

António Álvaro Pereira de Pinho

**Association of Human Papillomavirus with Cutaneous Squamous Cell  
Carcinoma and Basal Cell Carcinoma: a systematic review and meta-analysis**

março, 2018

António Álvaro Pereira de Pinho

Association of Human Papillomavirus with Cutaneous Squamous Cell Carcinoma  
and Basal Cell Carcinoma: a systematic review and meta-analysis

**Mestrado Integrado em Medicina**

**Área: Microbiologia Médica**

**Tipologia: Dissertação**

**Trabalho efetuado sob a Orientação de:**

**Professora Doutora Carmen Lisboa**

**E sob a Coorientação de:**

**Professor Doutor Alberto Freitas**

**Trabalho organizado de acordo com as normas da revista:**

**Journal of the European Academy of Dermatology and Venereology (JEADV)**

março, 2018

**FMUP**

Eu, António Álvaro Pereira de Pinho, abaixo assinado, nº mecanográfico 204202204, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 18/03/2018

Assinatura conforme cartão de identificação:

António Álvaro Pereira de Pinho

NOME

António Álvaro Pereira de Pinho

NÚMERO DE ESTUDANTE

20120201

E-MAIL

mimed12248@med.up.pt

DESIGNAÇÃO DA ÁREA DO PROJETO

Microbiologia Médica

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

Association of Human Papillomavirus with Cutaneous Squamous Cell Carcinoma and Basal Cell Carcinoma: a systematic review and meta-analysis

ORIENTADOR

Prof. Dra. Carmen Maria Lisboa da Silva

COORIENTADOR (se aplicável)

Prof. Dr. José Alberto da Silva Freitas

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 18/03/2018

Assinatura conforme cartão de identificação: António Álvaro Pereira de Pinho

*“Não sei o que nos espera mas sei o que me preocupa: é que a medicina, empolgada pela ciência, seduzida pela tecnologia e atordoada pela burocracia, apague a sua face humana e ignore a individualidade única de cada pessoa que sofre, pois embora se inventem cada vez mais modos de tratar, não se descobriu ainda a forma de aliviar o sofrimento sem empatia ou compaixão.”*

João Lobo Antunes

in "A Nova Medicina"

# **Association of Human Papillomavirus with Cutaneous Squamous Cell Carcinoma and Basal Cell Carcinoma: a systematic review and meta-analysis**

**Running Head:** Association of HPV with Cutaneous Squamous Cell Carcinoma and Basal Cell Carcinoma.

**Manuscript count:** 2967 words (body text, excluding figures, tables, legends, references, abstract and acknowledgment); 3 (three) tables; 6 (six) figures.

**Authors:** A.A. De-Pinho<sup>1</sup>, A.L.C. Ramalho<sup>2,3</sup>, J.V. Santos<sup>2,4,5</sup>, A. Freitas<sup>2,4</sup>, C. Lisboa<sup>1,4,6</sup>

<sup>1</sup> Division of Microbiology, Department of Pathology, Faculty of Medicine, University of Porto, Porto, Portugal

<sup>2</sup>MEDCIDS (Department of Community Medicine, Information and Health Decision Sciences), Faculty of Medicine, University of Porto, Portugal.

<sup>3</sup>FGV (Getúlio Vargas Foundation), Rio de Janeiro, Brazil

<sup>4</sup>CINTESIS (Center for Health Technology and Services Research), Porto, Portugal

<sup>5</sup>Public Health Unit, Aces Grande Porto VIII, Espinho/Gaia, Portugal

<sup>6</sup> Dermatovenereology Department, Centro Hospitalar S. João, EPE, Porto, Portugal

Corresponding author: António Álvaro Pereira de Pinho - mimed12248@med.up.pt;  
Alameda Professor Hernâni Monteiro 4200-319, Porto, Portugal; Fax: +351220426870.

**No funding sources.**

**No conflicts of interest.**

## **Abstract**

**Background** The association between human papillomavirus (HPV) and Non-Melanoma Skin Cancer (NMSC) remains undefined, with studies showing controversial results.

**Objective** We aim to determine if there is a significant association of HPV with Cutaneous Squamous Cell Carcinoma (cuSCC) and Basal Cell Carcinoma (BCC) and whether the prevalence of HPV is higher in those tumors from immunosuppressed patients when compared with tumors from immunocompetent patients.

**Methods** We conducted a systematic review and meta-analysis of observational studies. We searched four electronic databases up to March 2017, reviewed references of relevant articles and hand searched conference proceedings. The following search terms were used: skin cancer, squamous cell skin carcinoma, basal cell skin carcinoma and human papillomavirus. Only studies including biopsy samples evaluated by a pathologist, HPV detection by polymerase chain reaction (PCR) and a minimum of 10 cases and 10 controls were considered. Pooled effect size and 95% confidence intervals were calculated using random effects meta-analysis and the inverse variance method.

**Results** Of 2740 articles identified, 23 met the eligibility criteria. cuSCC were more likely to carry HPV than normal-appearing skin - pooled odds ratio (OR) 3.29 (95% CI 1.67-6.46). The overall association between HPV and BCC was also significant - pooled OR 2.56 (95% CI 1.48-5.14). An increase in HPV prevalence was found in cuSCC tumors from immunosuppressed patients when compared with immunocompetent patients - pooled OR 3.07 (95% CI 2.08-4.52). The same prevalence was not significant in BCC - pooled OR 2.38 (95% CI 0.95-5.96).

**Conclusion** These results support those studies that reported an association between HPV and NMSC, particularly in cuSCC and in tumors from immunosuppressed patients. Such evidence may have future research and clinical implications.

**Keywords** human papillomavirus; squamous skin cell carcinoma; basal skin cell carcinoma; meta-analysis; immunosuppression; systematic review.

## Introduction

Non-Melanoma Skin Cancer (NMSC) comprises Basal Cell Carcinoma (BCC) and Cutaneous Squamous Cell Carcinoma (cuSCC) as the major representatives.<sup>1</sup> BCC is the most common skin cancer<sup>2</sup> and is the most prevalent cancer in many countries worldwide.<sup>3</sup> On the other hand, cuSCC is the second most frequent form of skin cancer.<sup>4</sup> Reports from the World Health Organization estimate an occurrence of 2 to 3 million cases of NMSC annually worldwide.<sup>5</sup> The annual cost of treating skin cancers is estimated at \$8.1 billion, \$4.8 billion for NMSC in the United States.<sup>6</sup> NMSC is also a significant economic burden in European health expenditure, with €5.2 million annually costs for inpatient care attributable to these cancers in Sweden<sup>7</sup> and €105 to €130 million estimated annually costs of hospitalization in Germany.<sup>8</sup>

Ultraviolet Radiation (UVR) exposure is the main recognized risk factor for NMSC. About 90 percent of NMSC are associated with exposure to UVR from the sun<sup>9</sup>, but other risk factors have been studied as fair skin, older age and immunosuppression.<sup>10</sup> For instance, organ transplanted patients are approximately 100 times more likely to develop cuSCC than general population.<sup>11</sup> Considering the increased prevalence and the clinical behavior of NMSC and the high incidence of virally induced neoplasia such as Kaposi Sarcoma (Human Herpesvirus 8) in this cluster of patients, a viral etiology has also been hypothesized<sup>12, 13</sup> associated with NMSC.

More than 170 types of small capsid-enclosed double-stranded DNA human papillomaviruses (HPV) have been described.<sup>14</sup> Taxonomy recognizes five main HPV genera: alfa-, beta-, gamma-, mu- and nu- viruses.<sup>15</sup> Their oncogenic potential has been studied both *in vitro* and *in vivo* models demonstrating a well-established role of alfa-HPV in cervical carcinoma in an important portion of other anogenital, head and neck and oropharyngeal malignancies.<sup>16</sup>

In the last decades, the biology and contribution of HPV to NMSC development in previously healthy skin has been studied but controversial results were found and different methodological approaches were used.

Public health interventions have been launched to minimize the burden of HPV-associated malignancies, including worldwide vaccination programs against the most oncogenic viruses.<sup>16</sup> A better understanding of the relation between HPV and NMSC



would help to develop more effective preventive measures (such as vaccination) decreasing, therefore, the number of newly diagnosed cases and consequently the financial burden.

The main aim of the present study was to perform a systematic review and meta-analysis to determine the association of HPV with cuSCC and BCC. We also sought to ascertain whether the prevalence of HPV is higher in those tumors from immunosuppressed patients when compared with NMSC from immunocompetent patients.

## **Material and methods**

A systematic review and meta-analysis of observational studies focusing on the association between NMSC and the presence of human papillomaviruses in skin was performed. The methodology included definition of selection criteria and search strategy, quality assessment, data abstraction, bias risk assessment and statistical analysis. PRISMA statement was followed.<sup>17</sup>

### **Selection criteria**

The study eligibility criteria were determined before data collection in order to properly identify high quality studies appropriate for the analysis.

The following eligibility criteria were defined. Patient population: immunocompetent or immunosuppressed patients presenting healthy normal skin (controls) or non-anogenital NMSC lesions including Keratoacanthoma (KA), *in situ* cuSCC, invasive cuSCC or BCC (cases) reported by a pathologist, without other cutaneous comorbid conditions (Epidermolysis Verruciformis or other genodermatosis). Study design: observational studies with a minimum of 10 cases and 10 controls or a minimum of 20 patients. Exposure: HPV presence in skin detected by polymerase chain reaction-based (PCR) methods in biopsy samples. Outcome: NMSC occurrence in both immunocompetent and immunosuppressed subpopulations (Table 1). *INSERT T1*

### **Search strategy**

Our primary method to identify potentially eligible studies was an electronic biomedical literature search in the MEDLINE database, considering articles published until to March 10, 2017, written in English, using the following search keywords (skin cancer or squamous cell skin carcinoma or basal cell skin carcinoma) and human papillomavirus and MeSH terms: “skin neoplasms”, “carcinoma, squamous cell”, “carcinoma, basal cell” and “papillomaviridae”. Literature search was also performed, using the previous search keywords, in other three major databases: the Cochrane Library; the SciVerse Scopus; and the Web of Science.

We also reviewed the references list of all pertinent articles to recognize potentially appropriate studies. Abstracts from relevant conferences and scientific forums were also searched.

Duplicated articles were excluded and all the articles were uploaded to Covidence systematic review manager for quality assessment and further selection.

### **Quality assessment and data abstraction**

During the first phase of selection, each article was independently reviewed by two of the authors, based on its title and abstract to screen for relevance.

In the second phase, articles were independently full-text scrutinized by two reviewers, De-Pinho A.A. and Lisboa C. Selection criteria were applied, exclusions were decided and disagreements settled by consensus.

A standard data abstraction form was created by De-Pinho A.A. and Santos J.V. Each article was checked independently and all relevant data was extracted independently and in duplicate for each article by De-Pinho A.A. and Ramalho A.L.C. Any discrepancies in the duplicates were solved by the authors by discussing and achieving consensus.

General data extracted from each study included authors' names, year of publication, general study design, studied malignancies, tumour body locations, HPV subtypes that were searched, total numbers or HPV positive/HPV negative samples of cuSCC, BCC, normal skin, cuSCC or BCC in immunosuppressed or immunocompetent patients, depending on the hypothesis tested as well as other relevant information concerning methodological specificities to facilitate the assessment of bias risk. Extra data relative to age, sex, race, skin type, sun exposure, immunosuppression and PCR methodology details were also collected when available (Table 2). *INSERT T2*

Following the Newcastle-Ottawa Scale (NOS), the risk of bias in all reviewed studies was assessed regardless of the anticipated variability in outcomes or the validity of the included studies. Two reviewers performed the NOS assessment individually.

Evaluations were performed according to manual coding for observational studies, covering the domains: selection, comparability and exposure. Dispute resolutions between the two reviewers were conducted by consensus method.

### **Statistical methods**

For the assessment of the effect of HPV in the development of cuSCC or BCC, as well in the immunosuppression of both, we used the generic inverse variance method for both fixed and random effects estimation.

Generic inverse variance was used as data type for measuring the HPV effect in the outcomes as it is the optimal measure to describe retrospective exposure effects. Odds ratio (ORs) with the corresponding 95% confidence intervals (CI) were abstracted from each article. We identified the ORs reflecting the greatest degree of adjustment for possible confounding factors. When adjusted values were absent, we collected crude values. As a last resort, if ORs were not mentioned in the selected study, we calculated them based on the number of cases and controls described in the study. Random-effects method was used to pool the ORs and 95% CIs.

Heterogeneity of exposure effect was assessed by graphical inspection of forest plots and formally using the Q statistic (at a p value  $\leq 0.1$ ) and  $I^2$  statistic for studying studies' consistency.

Potential publication bias was assessed by visual analysis of funnel plots and analysis using Begg test.

Data processing and statistical analysis were performed using Review Manager (RevMan) software version 5.3.

## Results

### Search and study selection

The literature search protocol described above yielded a total of 2740 studies. During this process, two previous meta-analysis on the subject were identified – Wang et al<sup>18</sup> and Chahoud et al<sup>19</sup> and articles included in those studies were selected for our screening. No abstracts from relevant conferences and scientific forums were found.

After review of the titles and abstracts, a total of 2577 were excluded for non-compliance with the eligibility criteria. De-Pinho A.A. and Lisboa C. reviewed the full text of the remaining 163 articles. Selection criteria were applied, exclusions were decided and disagreements settled by consensus. After the final screening, 23 studies were selected (Fig. 1). *INSERT F1*

### Study characteristics

All the 23 included studies in our meta-analysis were case-controls (no cohort studies were identified). Depending on the outcome, a different number of studies was considered for statistical analysis (Fig. 1; Table 2). Some studies were included in more than one hypothesis analysis as they addressed more than one outcome.

Publication years ranged from 1996 to 2014. All studies included both male and female patients ranging all ages and resulted in a total of 558 cases and 907 controls for hypothesis A, 370 cases and 667 controls for hypothesis B, 444 cases and 286 controls for hypothesis C and 72 cases and 42 controls for hypothesis D.

After data collection, we recognized that, depending on the study, “cuSCC samples” included invasive cuSCC and/or in situ cuSCC and/or Keratoacanthoma. “Normal skin” was a reference to normal skin tissue from patients with cuSCC/BCC or normal skin samples from paired controls without tumours, depending on the studies, but always reported by a pathologist. Benign lesions were not included.

Some studies included in this meta-analysis evaluated several body locations, namely anogenital lesions.<sup>20</sup> Since those cases did not meet our eligibility criteria, they were excluded. This may explain a discrepancy between the values present in our meta-

analysis and the ones described in the original study. Samples that did not described the specific body location of the lesion or mentioned it only as “unknown” were included.

All the selected studies used PCR-based methods to identify the HPV presence in skin biopsies, but the genera/subtype detected was different among the case-controls, varying from broad spectrum, including alpha, beta and even gamma genera to only a specific HPV subtype.<sup>21</sup>

Only two of the selected studies presented adjusted OR: Forslund et al<sup>22</sup> – adjustment for age, sex, skin type, self-reported previous sunburns, eye colour and sun exposure; and Iftner et al<sup>23</sup> – adjustment for age, sex and sun exposure.

Additional information and abstracted data from the included studies are presented in Table 2.

#### **Methodological quality of included studies**

Quality assessment was performed using the NOS, covering the following domains: selection, comparability and exposure. Scores ranged from 5 to 9 (9 being the highest possible score), with a mean of 8.3 and median of 9 (Table 3).

Despite the presence of some possible bias in five studies<sup>23,24,25,26,27</sup> which scored 7 or less, they were relevant enough to be included. *INSERT T3*

#### **Publication bias**

Considering the existence of approximately symmetrical funnel plots among the studied hypothesis B, C and D, selected studies from these hypotheses seemed to show no evidence of publication bias. On the other hand, the funnel plot regarding hypothesis A showed some grade of asymmetry. For a better understanding, Begg test was applied to the four funnel plots and detected no evidence of publication bias among the four studied hypothesis (Fig. 2). *INSERT F2*

#### **Meta-analysis**

##### **Hypothesis A - HPV association with cuSCC *versus* normal skin**

A total of 11 studies comprising 558 cases and 907 controls were considered. In this pooled analysis, overall HPV-cuSCC association was significant with pooled OR 3.29

(CI 95%, 1.67-6.46,  $p < 0.001$ ), showing that squamous cell carcinomas were more likely to carry HPV than normal-appearing skin.  $I^2$  and Q statistics showed significant level of heterogeneity in the published studies, suggesting that individual study effect sizes varied based on different study designs,  $I^2 = 77.0\%$ ,  $Q = 43.94$ ,  $p < 0.001$  (Fig. 3). *INSERT F3*

#### **Hypothesis B - HPV association with BCC *versus* normal skin**

Seven studies met the eligibility criteria, resulting in a total of 370 cases and 667 controls. In our random effects pooled analysis, BCC was more likely to carry HPV than normal skin (pooled OR 2.56, CI 95%, 1.48-5.14,  $p = 0.008$ ).  $I^2$  and Q statistics showed moderate evidence of heterogeneity in the published studies,  $I^2 = 62.0\%$ ,  $Q = 15.82$ ,  $p = 0.01$  (Fig. 4). Removing studies one by one, we found that the exclusion of Caldeira et al<sup>21</sup>, considerable changed the level of heterogeneity. In the absence of this article, the pooled effect was OR 1.86, CI 95%, 1.10–3.14 and  $I^2 = 25.0\%$ ,  $p = 0.25$ . *INSERT F4*

#### **Hypothesis C - HPV association with cuSCC in immunosuppressed *versus* immunocompetent patients**

In this pooled analysis, 444 cases and 286 controls from 12 case-controls were included. cuSCC from immunosuppressed patients was significantly more likely to carry HPV than cuSCC samples from non-immunosuppressed patients (pooled OR 3.07, CI 95%, 2.08-4.52,  $p < 0.001$ ).  $I^2$  and Q values showed no significant level of heterogeneity in the published studies,  $I^2 = 22.0\%$ ,  $Q = 14.07$ ,  $p = 0.23$  (Fig. 5). *INSERT F5*

#### **Hypothesis D - HPV association with BCC in immunosuppressed *versus* immunocompetent patients**

Only 3 studies (72 cases and 42 controls) were included. The random effects pooled analysis on this outcome showed a non-statistically significant difference between immunosuppression and HPV presence in BCC (pooled OR 2.38, CI 95%, 0.95-5.96,  $p = 0.07$ ).  $I^2$  and Q statistics showed no significant level of heterogeneity in the published studies,  $I^2 = 0\%$ ,  $Q = 0.06$ ,  $p = 0.97$  (Fig. 6). *INSERT F6*

## Discussion

In our analysis studied populations varied from 23 samples (11 cases and 12 controls) to 540 samples (148 cases and 392 controls) with more than half of the studies including over 50 samples. About half of the selected case-controls were relatively recent (within the last 10 years). In general, the newer studies included broad-spectrum PCR techniques being able to detect more HPV types, specially beta-genera. Detected HPV varied from one specific HPV subtype to broad spectrum mucosal and cutaneous HPV, covering various genera.

Concerning cuSCC, these tumors were more likely to carry HPV than normal-appearing skin and immunosuppressed patients were more likely to carry HPV than immunocompetent patients. Therefore, our analysis shares similar results with the two previous identified meta-analysis on the subject.<sup>18,19</sup> However, Chahoud et al<sup>19</sup> did not included an analysis in immunosuppressed patients which represent a significant portion of the patients with cuSCC and allowed a wider range of methodologies to detect HPV, including multiplex and ELISA serologies.

Interestingly, HPV load seems to decrease during cuSCC carcinogenesis. Some studies refer a higher prevalence of HPV DNA in Actinic Keratosis (AK), a cuSCC premalignant lesion, comparing with the subsequent rates detected in cuSCC.<sup>28</sup> Another curious issue is that skin cells, unlike cells studied in the cervical region, do not require the expression of oncoproteins (like E6 and E7) by HPV viruses to preserve their transformed phenotype.<sup>28</sup> Finally, studies have failed to detect messenger RNA from beta-genus in cuSCC samples.<sup>29</sup> These three previous facts may suggest a temporary early role of HPV in the multistep process of oncogenesis and no need of these viruses for maintenance. We can speculate if HPV works as co-carcinogen factor with UVR. Actually, some studies have identified higher HPV rates in sun-exposed regions.<sup>22</sup> A local immunosuppression effect of UVR<sup>30</sup> or a direct trigger effect in viruses may induce apoptosis or DNA damage. This last theory was already demonstrated *in vitro* with cutaneous HPV types 5, 8, 20 and 77.<sup>31,32</sup>



There might also be different mechanisms driving tumor progression in HPV DNA positive versus HPV DNA negative cuSCC. Apart from the importance of studying the natural history of the carcinogenesis in HPV DNA positive cuSCC, immunological and genetic factors that are associated with HPV DNA negative tumors should also be better explored in the future.

Regarding BCC, no previous meta-analysis were found and the association with HPV was demonstrated for the first time in literature. Even without a statistical significance, BCC from immunosuppressed patients were more likely to carry HPV than tumors from immunocompetent patients.

Studies involving BCC tumours were, in general, more recent<sup>21-23, 25, 33-38</sup> and a lack in both quantity and quality of case-controls on BCC, particularly involving immunosuppressed patients (low number of cases and controls for hypothesis D) was found. This fact may be due to an easier logical reasoning relating cuSCC with HPV (rather than BCC), as the causality of papillomaviruses and carcinoma of the cervix in women (affecting squamous cells) is well-established. Additionally, in contrast to the immunocompetent population, cuSCC is more common in immunosuppression context than BCC.<sup>39</sup> New studies, covering a larger number of patients with BCC and preferentially prospective designed, would improve the quality of future meta-analysis in this matter.

The role of HPV in cutaneous carcinogenesis is far away from being completely understood. Even though our results demonstrate a strong association between HPV and NMSC, a higher prevalence of the virus in tumoral tissues does not imply causality. It is of great importance to differentiate if the HPV infection is only a simultaneously phenomenon with the occurrence of neoplasia or a major agent during the process of carcinogenesis.<sup>40</sup> Thus, cohort studies should be designed and performed to assess this causality, adjusting for other risk factors.

Definition of a causality link between NMSC and HPV may lead to new diagnostic and therapeutic tools. For instance, the well-established relationship between cervical carcinoma and HPV provided effective HPV vaccines and a better control of this cancer.<sup>16</sup> Acquired immunity through vaccination against selected HPV subtypes could potentially

be explored as an effective preventive approach, decreasing NMSC burden and further costs of care.

This meta-analysis has some limitations that should be pointed out.

Unfortunately, most of the studies did not present adjusted measures for possible confounding factors, particularly for exposure to UVR and older age, known NMSC risk factors.

Other important limitation is the great degree of heterogeneity among the literature regarding sampling methods, the laboratory technique to HPV detection and the HPV type studied. We specifically tried to minimize these points, considering only studies based on biopsy samples evaluated by experienced pathologists or PCR techniques for HPV detection rather than eyebrow pluck, skin swab or serologies.

### **Conclusions**

This study represents, to our knowledge, the most extensive meta-analysis assessing the epidemiological association of HPV with cuSCC using PCR analysis.

Notably, this is the first meta-analysis evaluating the association between HPV and BCC. This work provides additional evidence of the eventual involvement of HPV in the development of these tumors in both immunocompetent and immunosuppressed individuals.

### **Acknowledgments**

We thank the precious help of Luís F. Azevedo, MD PhD, in the invaluable advices on the pooled analysis.

## References:

1. Eisemann N, Waldmann A, Geller AC, Weinstock MA, Volkmer B, Greinert R, et al. Non-melanoma skin cancer incidence and impact of skin cancer screening on incidence. *J Invest Dermatol*. 2014;**134**(1):43-50.
2. Mohan SV, Chang AL. Advanced Basal Cell Carcinoma: Epidemiology and Therapeutic Innovations. *Curr Dermatol Rep*. 2014;**3**:40-5.
3. Lomas A, Leonardi-Bee J, Bath-Hextall F. A systematic review of worldwide incidence of nonmelanoma skin cancer. *Br J Dermatol*. 2012;**166**(5):1069-80.
4. Karia PS, Han J, Schmults CD. Cutaneous squamous cell carcinoma: estimated incidence of disease, nodal metastasis, and deaths from disease in the United States, 2012. *J Am Acad Dermatol*. 2013;**68**(6):957-66.
5. WHO - World Health Organization. *Skin Cancers*. 2017. Available at: <http://www.who.int/uv/faq/skincancer/en/index1.html> (last accessed 15 March 2017).
6. Guy GP, Jr., Machlin SR, Ekwueme DU, Yabroff KR. Prevalence and costs of skin cancer treatment in the U.S., 2002-2006 and 2007-2011. *Am J Prev Med*. 2015;**48**(2):183-7.
7. Tinghog G, Carlsson P, Synnerstad I, Rosdahl I. Societal cost of skin cancer in Sweden in 2005. *Acta Derm Venereol*. 2008;**88**(5):467-73.
8. Stang A, Stausberg J, Boedeker W, Kerek-Bodden H, Jockel KH. Nationwide hospitalization costs of skin melanoma and non-melanoma skin cancer in Germany. *J Eur Acad Dermatol Venereol*. 2008;**22**(1):65-72.
9. Koh HK, Geller AC, Miller DR, Grossbart TA, Lew RA. Prevention and early detection strategies for melanoma and skin cancer. Current status. *Arch Dermatol*. 1996;**132**(4):436-43.
10. Euvrard S, Kanitakis J, Claudy A. Skin cancers after organ transplantation. *N Engl J Med*. 2003;**348**(17):1681-91.
11. Lindelof B, Sigurgeirsson B, Gabel H, Stern RS. Incidence of skin cancer in 5356 patients following organ transplantation. *Br J Dermatol*. 2000;**143**(3):513-9.
12. Feltkamp MC, de Koning MN, Bavinck JN, Ter Schegget J. Betapapillomaviruses: innocent bystanders or causes of skin cancer. *J Clin Virol*. 2008;**43**(4):353-60.
13. Vajdic CM, McDonald SP, McCredie ME, et al. Cancer incidence before and after kidney transplantation. *JAMA*. 2006;**296**(23):2823-31.
14. de Villiers EM. Cross-roads in the classification of papillomaviruses. *Viol J*. 2013;**445**(1-2):2-10.
15. Bernard HU, Burk RD, Chen Z, van Doorslaer K, zur Hausen H, de Villiers EM. Classification of papillomaviruses (PVs) based on 189 PV types and proposal of taxonomic amendments. *Viol J*. 2010;**401**(1):70-9.
16. Grce M, Mravak-Stipetic M. Human papillomavirus-associated diseases. *Clin Dermatol*. 2014;**32**(2):253-8.
17. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009;**339**:b2700.
18. Wang J, Aldabagh B, Yu J, Arron ST. Role of human papillomavirus in cutaneous squamous cell carcinoma: a meta-analysis. *J Am Acad Dermatol*. 2014;**70**(4):621-9.
19. Chahoud J, Semaan A, Chen Y, Cao M, Rieber AG, Rady P, et al. Association Between beta-Genus Human Papillomavirus and Cutaneous Squamous Cell Carcinoma

in Immunocompetent Individuals-A Meta-analysis. *JAMA Dermatol.* 2016;**152**(12):1354-64.

20. Hama N, Ohtsuka T, Yamazaki S. Detection of mucosal human papilloma virus DNA in bowenoid papulosis, Bowen's disease and squamous cell carcinoma of the skin. *J Dermatol.* 2006;**33**(5):331-7.

21. Caldeira S, Zehbe I, Accardi R, Malanchi I, Dong W, Giarre M, et al. The E6 and E7 proteins of the cutaneous human papillomavirus type 38 display transforming properties. *J Virol.* 2003;**77**(3):2195-206.

22. Forslund O, Iftner T, Andersson K, Lindelof B, Hradil E, Nordin P, et al. Cutaneous human papillomaviruses found in sun-exposed skin: Beta-papillomavirus species 2 predominates in squamous cell carcinoma. *J Infect Dis.* 2007;**196**(6):876-83.

23. Iftner A, Klug SJ, Garbe C, Blum A, Stancu A, Wilczynski SP, et al. The prevalence of human papillomavirus genotypes in nonmelanoma skin cancers of nonimmunosuppressed individuals identifies high-risk genital types as possible risk factors. *Cancer Res.* 2003;**63**(21):7515-9.

24. Asgari MM, Kiviat NB, Critchlow CW, Stern JE, Argenyi ZB, Raugi GJ, et al. Detection of human papillomavirus DNA in cutaneous squamous cell carcinoma among immunocompetent individuals. *J Invest Dermatol.* 2008;**128**(6):1409-17.

25. Bernat-Garcia J, Morales Suarez-Varela M, Vilata-Corell JJ, Marquina-Vila A. Detection of human papillomavirus in nonmelanoma skin cancer lesions and healthy perilesional skin in kidney transplant recipients and immunocompetent patients. *Actas Dermosifiliogr.* 2014;**105**(3):286-94.

26. Mackintosh LJ, de Koning MN, Quint WG, Ter Schegget J, Morgan IM, Herd RM, et al. Presence of beta human papillomaviruses in nonmelanoma skin cancer from organ transplant recipients and immunocompetent patients in the West of Scotland. *Br J Dermatol.* 2009;**161**(1):56-62.

27. Plasmeijer EI, Neale RE, Buettner PG, de Koning MNC, ter Schegget J, Quint WGV, et al. Betapapillomavirus infection profiles in tissue sets from cutaneous squamous cell-carcinoma patients. *Int J Cancer.* 2010;**126**(11):2614-21.

28. Weissenborn SJ, Nindl I, Purdie K, Harwood C, Proby C, Breuer J, et al. Human papillomavirus-DNA loads in actinic keratoses exceed those in non-melanoma skin cancers. *J Invest Dermatol.* 2005;**125**(1):93-7.

29. Arron ST, Ruby JG, Dybbro E, Ganem D, Derisi JL. Transcriptome sequencing demonstrates that human papillomavirus is not active in cutaneous squamous cell carcinoma. *J Invest Dermatol.* 2011;**131**(8):1745-53.

30. Kripke ML. Ultraviolet radiation and immunology: something new under the sun-presidential address. *Clin Cancer Res.* 1994;**54**(23):6102-5.

31. Purdie KJ, Pennington J, Proby CM, Khalaf S, de Villiers EM, Leigh IM, et al. The promoter of a novel human papillomavirus (HPV77) associated with skin cancer displays UV responsiveness, which is mediated through a consensus p53 binding sequence. *EMBO J.* 1999;**18**(19):5359-69.

32. Akgul B, Lemme W, Garcia-Escudero R, Storey A, Pfister HJ. UV-B irradiation stimulates the promoter activity of the high-risk, cutaneous human papillomavirus 5 and 8 in primary keratinocytes. *Arch Virol.* 2005;**150**(1):145-51.

33. Berkhout RJ, Bouwes Bavinck JN, ter Schegget J. Persistence of human papillomavirus DNA in benign and (pre)malignant skin lesions from renal transplant recipients. *Eur J Clin Microbiol Infect Dis.* 2000;**38**(6):2087-96.

34. Escutia B, Ledesma E, Serra-Guillen C, Gimeno C, Vilata JJ, Guillen C, et al. Detection of human papilloma virus in normal skin and in superficial and nodular basal cell carcinomas in immunocompetent subjects. *J Eur Acad Dermatol Venereol*. 2011;**25**(7):832-8.
35. Harwood CA, Suretheran T, McGregor JM, Spink PJ, Leigh IM, Breuer J, et al. Human papillomavirus infection and non-melanoma skin cancer in immunosuppressed and immunocompetent individuals. *J Med Virol*. 2000;**61**(3):289-97.
36. Reuschenbach M, Tran T, Faulstich F, Hartschuh W, Vinokurova S, Kloor M, et al. High-risk human papillomavirus in non-melanoma skin lesions from renal allograft recipients and immunocompetent patients. *Br J Cancer*. 2011;**104**(8):1334-41.
37. Rollison DE, Pawlita M, Giuliano AR, Iannacone MR, Sondak VK, Messina JL, et al. Measures of cutaneous human papillomavirus infection in normal tissues as biomarkers of HPV in corresponding nonmelanoma skin cancers. *Int J Cancer*. 2008;**123**(10):2337-42.
38. Zaravinos A, Kanellou P, Spandidos DA. Viral DNA detection and RAS mutations in actinic keratosis and nonmelanoma skin cancers. *Br J Dermatol*. 2010;**162**(2):325-31.
39. Ramsay HM, Fryer AA, Reece S, Smith AG, Harden PN. Clinical risk factors associated with nonmelanoma skin cancer in renal transplant recipients. *Am J Kidney Dis*. 2000;**36**(1):167-76.
40. Rothman KJ, Greenland S. Causation and causal inference in epidemiology. *Am J Public Health*. 2005;**95** Suppl 1:S144-50.
41. Arends MJ, Benton EC, McLaren KM, Stark LA, Hunter JA, Bird CC. Renal allograft recipients with high susceptibility to cutaneous malignancy have an increased prevalence of human papillomavirus DNA in skin tumours and a greater risk of anogenital malignancy. *Br J Cancer*. 1997;**75**(5):722-8.
42. Cairey-Remonnay S, Humbey O, Mougin C, Algros MP, Mauny F, Kanitakis J, et al. TP53 Polymorphism of Exon 4 at Codon 72 in Cutaneous Squamous Cell Carcinoma and Benign Epithelial Lesions of Renal Transplant Recipients and Immunocompetent Individuals: Lack of Correlation with Human Papillomavirus Status. *J Invest Dermatol*. 2002;**118**(6):1026-31.
43. Forslund O, Ly H, Reid C, Higgins G. A broad spectrum of human papillomavirus types is present in the skin of Australian patients with non-melanoma skin cancers and solar keratosis. *Br J Dermatol*. 2003;**149**(1):64-73.
44. Forslund O, DeAngelis PM, Beigi M, Schjolberg AR, Clausen OP. Identification of human papillomavirus in keratoacanthomas. *J Cutan Pathol*. 2003;**30**(7):423-9.
45. Gustafsson AC, Ren ZP, Asplund A, Ponten F, Lundeberg J. The role of p53 codon 72 and human papilloma virus status of cutaneous squamous cell carcinoma in the Swedish population. *Acta Derm Venereol*. 2004;**84**(6):439-44.
46. Purdie KJ, Suretheran T, Sterling JC, Bell L, McGregor JM, Proby CM, et al. Human papillomavirus gene expression in cutaneous squamous cell carcinomas from immunosuppressed and immunocompetent individuals. *J Invest Dermatol*. 2005;**125**(1):98-107.
47. Shamanin V, zur Hausen H, Lavergne D, Proby CM, Leigh IM, Neumann C, et al. Human papillomavirus infections in nonmelanoma skin cancers from renal transplant recipients and nonimmunosuppressed patients. *J Natl Cancer Inst*. 1996;**88**(12):802-11.

48. Stockfleth E, Nindl I, Sterry W, Ulrich C, Schmook T, Meyer T. Human papillomaviruses in transplant-associated skin cancers. *Dermatol Surg*. 2004;**30**(4 Pt 2):604-9.

## Tables

Table 1

---

### Eligibility Criteria

---

Article in **English, excluding abstracts, reviews and letters to the editor**

Minimum **of 10 cases and 10 controls/20 patients** for determination of Odds Ratio (OR)

Patients **without other cutaneous comorbid conditions** (Epidermodysplasia Verruciformis or other genodermatosis)

Samples including **all skin body areas except anogenital**

**Biopsy samples only** (rather than eyebrow pluck, skin swab or serology)

**HPV detection by PCR-based methods**

**Outcome** (at least one):

- For hypothesis A – **HPV-PCR positivity in cuSCC *versus* normal skin**
  - For hypothesis B – **HPV-PCR positivity in BCC *versus* normal skin**
  - For hypothesis C – **HPV-PCR positivity in cuSCC from immunosuppressed patients *versus* cuSCC from immunocompetent patients**
  - For hypothesis D – **HPV-PCR positivity in BCC from immunosuppressed patients *versus* BCC from immunocompetent patients**
-



Table 2

Source	Design and population (hypothesis)	HPV subtypes	Malignancies	Tumor Body Locations
Arends et al, <sup>41</sup> 1997	Case-control 27 cases 15 controls – C	HPV subtypes 1, 2, 5, 8, 6b, 11, 16 and 18	cuSCC	Sun-exposed and non-sun-exposed areas
Arron et al, <sup>29</sup> 2011	Case-control 67 cases 18 controls – A 39 cases 28 controls – C	Broad spectrum beta-HPV	cuSCC, KA type included	Sun-exposed and non-sun-exposed areas
Asgari et al, <sup>24</sup> 2008	Case-control 85 cases 190 controls – A	Broad spectrum HPV (including alpha, beta and gamma genera)	cuSCC	Sun-exposed and non-sun-exposed areas
Berkhout et al, <sup>33</sup> 2000	Case-control 81 cases 31 controls – A 14 cases 31 controls – B	Broad spectrum HPV (including alpha and beta genera)	cuSCC and BCC	Sun-exposed and non-sun-exposed areas
Bernat-García et al, <sup>25</sup> 2014	Case-control 17 cases 17 controls – C 13 cases 13 controls – D	Broad spectrum HPV (including alpha, beta and gamma genera)	cuSCC and BCC	Sun-exposed and non-sun-exposed areas
Cairey-Remonnay et al, <sup>42</sup> 2002	Case-control 53 cases 51 controls – C	Broad spectrum of mucosal and cutaneous HPV	cuSCC	Sun-exposed areas
Caldeira et al, <sup>21</sup> 2003	Case-control 26 cases 41 controls – A 69 cases 41 controls – B	HPV subtype 38	cuSCC and BCC	ND
Escutia et al, <sup>34</sup> 2011	Case-control 70 cases 29 controls – B	Broad spectrum HPV (including alpha, beta and gamma genera)	BCC	Sun-exposed and non-sun-exposed areas
Forslund et al, <sup>43</sup> 2003	Case-control 11 cases 12 controls – C	Broad spectrum of cutaneous HPV	cuSCC	Sun-exposed and non-sun-exposed areas
Forslund et al, <sup>44</sup> 2003, 2nd article	Case-control 60 cases 12 controls – C	Broad spectrum of cutaneous HPV	cuSCC, only KA	ND
Forslund et al, <sup>22</sup> 2007	Case-control 82 cases 392 controls – A 148 cases 392 controls – B	Broad spectrum HPV (including alpha, beta and gamma genera)	cuSCC and BCC	Sun-exposed and non-sun-exposed areas
Gustafsson et al, <sup>45</sup> 2004	Case-control 36 cases 27 controls – A	Broad anogenital HPV	cuSCC	ND
Hama et al, <sup>20</sup> 2006	Case-control 23 cases 17 controls – A	Broad spectrum of mucosal HPV	cuSCC	Sun-exposed and non-sun-exposed areas
Harwood et al, <sup>35</sup> 2000	Case-control 44 cases 22 controls – C 24 cases 11 controls – D	Broad spectrum of cutaneous, mucosal and EV HPV	cuSCC and BCC	Sun-exposed and non-sun-exposed areas
Iftner et al, <sup>23</sup> 2003	Case-control 72 cases 106 controls – A 18 cases 106 controls – B	Broad spectrum of mucosal and EV HPV	cuSCC and BCC	Sun-exposed and non-sun-exposed areas
Mackintosh et al, <sup>26</sup> 2009	Case-control 53 cases 11 controls – A 30 cases 23 controls – C	Broad spectrum beta-HPV	cuSCC, KA included in IS group	ND
Plasmeijer et al, <sup>27</sup> 2010	Case-control 21 cases 21 controls – A	Broad spectrum beta-HPV	cuSCC	Sun-exposed and non-sun-exposed areas
Purdie et al, <sup>46</sup> 2005	Case-control 84 cases 17 controls – C	Broad spectrum cutaneous HPV	cuSCC	ND
Reuschenbach et al, <sup>36</sup> 2011	Case-control 43 cases 44 controls – C 35 cases 18 controls – D	HPV subtypes 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68, 73, and 82	cuSCC, KA included and BCC	Sun-exposed and non-sun-exposed areas
Rollison et al, <sup>37</sup> 2008	Case-control 13 cases 15 controls – B	Broad spectrum beta-HPV	BCC	Sun-exposed and non-sun-exposed areas
Shamanin et al, <sup>47</sup> 1996	Case-control 20 cases 26 controls – C	Broad spectrum HPV	cuSCC	ND
Stockfleth et al, <sup>48</sup> 2004	Case-control 16 cases 19 controls – C	Broad spectrum mucosal, EV and classical wart-associated HPV	cuSCC	ND
Zaravinos et al, <sup>38</sup> 2010	Case-control 12 cases 53 controls – A 38 cases 53 controls – B	Broad spectrum HPV	cuSCC and BCC	Sun-exposed and non-sun-exposed areas

Table 3

Source	Newcastle-Ottawa Score	OR Adjustment
Arends et al, <sup>41</sup> 1997	8	--
Arron et al, <sup>29</sup> 2011	8	--
Asgari et al, <sup>24</sup> 2008	7	--
Berkhout et al, <sup>33</sup> 2000	9	--
Bernat-García et al, <sup>25</sup> 2014	7	--
Cairey-Remonnay et al, <sup>42</sup> 2012	9	--
Caldeira et al, <sup>21</sup> 2003	9	--
Escutia et al, <sup>34</sup> 2010	9	--
Forslund et al, <sup>43</sup> 2003	9	--
Forslund et al, <sup>44</sup> 2003, 2nd article	9	--
Forslund et al, <sup>22</sup> 2007	9	Adjusted for age, sex, skin type, self-reported previous sunburns, eye color and sun exposure at biopsy site
Gustafson et al, <sup>45</sup> 2004	9	--
Hama et al, <sup>20</sup> 2006	9	--
Harwood et al, <sup>35</sup> 2000	9	--
Iftner et al, <sup>23</sup> 2003	5	Adjusted for age, sex and sun exposure (sun exposed: head, face, neck, forearm, hands, and lower limb)
Mackintosh et al, <sup>26</sup> 2009	5	--
Plasmeijer et al, <sup>27</sup> 2010	7	--
Purdie et al, <sup>46</sup> 2005	9	--
Reuschenbach et al, <sup>36</sup> 2011	9	--
Rollison et al, <sup>37</sup> 2008	9	--
Shamanin et al, <sup>47</sup> 1996	9	--
Stockfleth et al, <sup>48</sup> 2004	9	--
Zaravinos et al, <sup>38</sup> 2010	9	--

## Figures

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6





## Legends

**Table 1** – Eligibility criteria for systematic review.

**Table 2** – Characteristics of studies meeting search eligibility criteria.

IS – Immunosuppressed; KA – Keratoacanthoma; ND – Not defined.

**Table 3** - Newcastle-Ottawa scale for assessment of quality and OR adjustment of the included studies.

**Fig. 1** - Flow chart of search strategy for the meta-analysis and number of studies included in final quantitative analysis by hypothesis.

\* Records included in previous meta-analysis and that were not selected via research protocol.

\*\* Some studies were included in more than one hypothesis analysis as they met criteria for both.

**Fig. 2** – Funnel plots with effect measures (Odds Ratio (OR) as a function of its standard error (SE)) for the outcome of HPV positivity in studies comparing (a) cuSCC versus normal skin – Begg test:  $p = 0.2183$ ; (b) BCC versus normal skin – Begg test:  $p = 0.8406$ ; (c) cuSCC from immunosuppressed patients versus cuSCC from immunocompetent patients – Begg test:  $p = 0.7726$ ; (d) BCC from immunosuppressed patients versus BCC from immunocompetent patients – Begg test:  $p = 1.0000$ .

**Fig. 3** – Forest plot with pooled effect size and 95% confidence intervals (CI) for hypothesis A - HPV-PCR positivity in cuSCC *versus* normal skin. The squares and horizontal lines correspond to the study-specific odds ratios (ORs) and 95% CIs. The diamond shape represents the pooled OR and 95% CI of the overall population.

**Fig. 4** – Forest plot with pooled effect size and 95% confidence intervals (CI) for hypothesis B - HPV-PCR positivity in BCC *versus* normal skin. The squares and horizontal lines correspond to the study-specific odds ratios (ORs) and 95% CIs. The diamond shape represents the pooled OR and 95% CI of the overall population.

**Fig. 5** – Forest plot with pooled effect size and 95% confidence intervals (CI) for hypothesis C - HPV-PCR positivity in cuSCC from immunosuppressed patients *versus* cuSCC from immunocompetent patients. The squares and horizontal lines correspond to the study-specific odds ratios (ORs) and 95% CIs. The diamond shape represents the pooled OR and 95% CI of the overall population.

**Fig. 6** – Forest plot with pooled effect size and 95% confidence intervals (CI) for hypothesis D - HPV-PCR positivity in BCC from immunosuppressed patients *versus* BCC from immunocompetent patients. The squares and horizontal lines correspond to the study-specific odds ratios (ORs) and 95% CIs. The diamond shape represents the pooled OR and 95% CI of the overall population.

## **Agradecimentos**

À minha orientadora, Professora Doutora Carmen Lisboa, por ter aceite o desafio de me orientar, pela disponibilidade inestimável e pelo constante *role model* de docência e profissionalismo ao longo dos últimos anos.

Ao meu orientador, Professor Doutor Alberto Freitas, ao Dr. André Ramalho e ao Dr. João Vasco Santos, pela contribuição científica, sugestões e correções que tanto engradeceram a qualidade deste trabalho “de equipa”.

À minha família, amigos, colegas de curso e companheiros de residência que marcaram (e marcam todos os dias) este meu percurso orgulhosamente trilhado.

## **Anexo - Normas de Publicação da Revista**

### ***Journal of the European Academy of Dermatology and Venereology (JEADV)***

Editor-in-Chief: Prof. Dr. med. Dr. phil. Johannes Ring, München, Germany

Impact Factor: 3.528

ISI Journal Citation Reports © Ranking: 2016: 9/63 (Dermatology)

Online ISSN: 1468-3083

## **1. AIMS & SCOPE**

The *Journal of the European Academy of Dermatology and Venereology (JEADV)* is the official organ of the European Academy of Dermatology and Venereology (EADV).

*JEADV* publishes articles of general and practical interest in the field of dermatology and venereology including clinical and basic science topics, as well as research with practical implications. It does so through editorials, review and practice articles, original papers of general interest, short reports, case reports, letters to the editor, news items, features and Academy announcements.

## **2. MANUSCRIPT CATEGORIES**

*JEADV* invites the following types of submissions:

**Original articles.** Original articles are the Journal's primary mode of communication. Original articles must include a structured abstract (maximum 300 words), and should not exceed 3000 words of body text.

(Manuscripts reporting randomised controlled trials (RCTs) must follow the **CONSORT** statement. RCTs will not be considered by *JEADV* without submission of a completed **CONSORT checklist**.)

**Review articles.** The Journal is particularly keen to publish concise, high-quality review articles of recent advances in laboratory or clinical research. Review articles may be solicited by the Editor-in-Chief or may be submitted by authors for publication subject to peer review. Review articles must include an unstructured abstract (maximum 300 words), and should not exceed 3000 words of body text. Use of tables and figures is encouraged.

**Short reports.** Brief data papers are published as short reports. Short reports must include a structured abstract and should not exceed 1500 words of body text, 4 figures/tables and 20 references.

**Letters to the Editor.** Letters to the Editor (Correspondence) may be in response to issues arising from recently published articles, or short, free-standing pieces expressing an opinion. Case reports will be considered, but published as items of correspondence. Letters to the Editor do not require abstracts, should be formatted in one continuous section and should not exceed 600 words, 10 references and two figures. All letters are subject to expert review.

### **3. SUBMISSION OF MANUSCRIPTS**

All submissions should be made online at the *JEADV* ScholarOne Manuscripts (formerly known as Manuscript Central) site—<http://mc.manuscriptcentral.com/jeadv>

New users should first create an account. Once a user is logged onto the site, submissions should be made via the Author Centre. Authors must also supply:

(i) completed **Conflicts of Interest Disclosure form(s)**—Each listed author must complete the form electronically. It is the responsibility of the corresponding author to upload—on behalf of all co-authors—completed forms as ‘Supplementary Material for Review’ via ScholarOne Manuscripts (formerly known as Manuscript Central) at the same time as manuscript submission. Manuscripts will not be sent for peer-review without these forms. (See section 6 for further information.)

#### 4. CONTACTING THE EDITORIAL OFFICE

**Asao Sarukawa**, *JEADV* Editorial Office, EADV Headquarters, Via S. Balestra 22 B. 6900 Lugano, Switzerland

Tel: +41 91 973 45 20; +41 91 973 45 30; Email: [jeadv@eadv.org](mailto:jeadv@eadv.org)

#### 5. PREPARATION OF MANUSCRIPTS

Manuscripts must be written in English.

Text should be supplied in a format compatible with Microsoft Word for Windows (PC). Charts and tables are considered textual and should also be supplied in a format compatible with Word. All figures (illustrations, diagrams, photographs) should be supplied in jpg, tiff or eps format.

All manuscripts must be typed in 12 pt font with margins of at least 2.5 cm. Submissions must comply with the word limits defined in section 2 and, where appropriate, include:

**Title page.** The first page of all manuscripts should contain the following information:

1. the title of the paper
2. a running head not exceeding 50 characters
3. manuscript word, table and figure count
4. names of authors as initial(s) followed by surnames
5. names of the institutions at which the research was conducted, clearly linked to respective authors
6. name, address, telephone and fax number, and email address of corresponding author
7. a statement of all funding sources that supported the work
8. any conflict of interest disclosures (see sections 3 and 6).

**Abstracts.** Authors submitting original articles should note that structured abstracts are required. The structured abstract should adopt the format: Background, Objective, Methods, Results, Conclusion.

Review articles require abstracts but they need not be structured. Letters to the Editor do not require abstracts. Abstracts should contain no citations to previously published work.

**Text.** This should in general, but not necessarily, be divided into sections with the headings: Introduction, Materials and Methods, Results, Discussion, Acknowledgments, References, Tables, Legends and Figures.

**Tables and figures.** Tables and figures should not be inserted in the appropriate place in the text but should be included at the end of the manuscript, each on a separate page. Tables and figures should be referred to in text as follows: Fig. 1, Figs 2–4; Table 1, Table 2. The place at which a table or figure is to be inserted in the printed text should be indicated clearly on the manuscript. Each table and/or figure must have a legend that explains its purpose without reference to the text. Where a figure has more than one panel, each panel should be labelled in the top left-hand corner using lower case letters in parentheses, i.e., (a), (b), etc., and a brief description of each panel given in the figure legend.

Colour illustrations are welcomed and all colour is published free of charge to the author.

Authors are themselves responsible for obtaining permission to reproduce previously published figures or tables. When an individual is identifiable in a photograph written permission must be obtained (see section 6).

If submitting artwork electronically, please read the information at: <http://authorservices.wiley.com/bauthor/illustration.asp>

**References.** References should be in Vancouver format and appear as consecutive, unbracketed superscript numbers in the text, e.g., ‘in our previous reports<sup>1,2</sup> and those of Smith *et al.*<sup>3–5</sup>’ and should be listed numerically in the reference list at the end of the article.

Format references as below, using standard (Medline) abbreviations for journal titles. If more than six authors, include the first three authors followed by *et al.* If six or fewer authors, please include all authors' names.

1. de Berker DAR, Baran R, Dawber RPR. The nail in dermatological diseases. In: *Baran and Dawber's Diseases of the Nails and their Management* (Baran R, Dawber RPR, de Berker DAR, Haneke E, Tosti A, eds), 3rd edn. Oxford: Blackwell Science Ltd, 2001; 172–92.
2. Wollina U, Hansel G. The use of topical calcineurin inhibitors in lupus erythematosus: an overview. *J Eur Acad Dermatol Venereol* 2008;**22**:1–6.
3. Graham-Brown R, Burns T. *Lecture Notes: Dermatology*. Oxford: Wiley-Blackwell, 2006.
4. British Lymphology Society. *Consensus Document on the Management of Cellulitis in Lymphoedema*. 2007. Available at: [http://www.lymphoedema.org/lsn/consensus\\_on\\_cellulitis\\_dec\\_06.pdf](http://www.lymphoedema.org/lsn/consensus_on_cellulitis_dec_06.pdf) (last accessed 28 November 2007).

**Reporting standards.** Manuscripts reporting randomised controlled trials (RCTs) must follow the **CONSORT** statement. RCTs will not be considered by *JEADV* without submission of a completed **CONSORT checklist**. Checklists should be uploaded during manuscript submission using file designation 'Supplementary files for review'.

## 6. DECLARATIONS

**Original publication.** Submission of a manuscript will be held to imply that it contains original unpublished work and is not being submitted for publication elsewhere at the same time. The author must supply a full statement to the Editor about all submissions and previous reports that might be regarded as redundant or duplicate publication of the same or very similar work.

**Conflicts of interest.** Authors are responsible for disclosing all financial and personal relationships between themselves and others that might be perceived by others as



biasing their work. To prevent ambiguity, authors must state explicitly whether potential conflicts do or do not exist. (See section 3.)

**Ethics.** When reporting experiments on human subjects, indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional or regional) and with the Helsinki Declaration of 1975, as revised in 1983. Do not use patients' names, initials or hospital numbers, especially in illustrative material. When reporting experiments on animals, indicate whether the institution's or a national research council's guide for, or any national law on, the care and use of laboratory animals was followed. A statement describing explicitly the ethical background to the studies being reported should be included in all manuscripts in the Materials and Methods section. Ethics committee or institutional review board approval should be stated.

Patients have a right to privacy that should not be infringed without informed consent. Identifying information should not be published in written descriptions, photographs and pedigrees unless the information is essential for scientific purposes and the patient (or parent or guardian) gives written informed consent for publication. Identifying details should be omitted if they are not essential but patient data should never be altered or falsified in an attempt to attain anonymity. Complete anonymity is difficult to achieve and informed consent should be obtained if there is any doubt. For example, masking the eye region in photographs of patients is inadequate protection of anonymity.

**Authorship.** All persons designated as authors should qualify for authorship and all those who qualify should be listed. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content. One or more authors should take responsibility for the integrity of the work as a whole, from inception to published article. Authorship credit should be based only on (i) substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; (ii) drafting the article or revising it critically for important intellectual content; (iii) final approval of the version to be published. Conditions (i), (ii) and (iii) must all be met. Acquisition of funding, the collection of data or general

supervision of the research group, by themselves, do not justify authorship. All others who contributed to the work who are not authors should be named in the Acknowledgments section.

**Committee on Publication Ethics (COPE).** As a member of the Committee on Publication Ethics (COPE), adherence to these submission criteria is considered essential for publication in the *JEADV*; mandatory fields are included in the online submission process to ensure this. If, at a later stage in the submission process or even after publication, a manuscript or authors are found to have disregarded these criteria, it is the duty of the Editor to report this to COPE. COPE may recommend that action be taken, including but not exclusive to, informing the authors' professional regulatory body and/or institution of such a dereliction. The website for COPE may be accessed at: <http://www.publicationethics.org.uk>

## **7. ADDITIONAL INFORMATION ON ACCEPTANCE**

**Copyright.** If your paper is accepted, the author identified as the formal corresponding author for the paper will receive an email prompting them to login into Author Services; where via the Wiley Author Licensing Service (WALS) they will be able to complete the license agreement on behalf of all authors on the paper.

### **For authors signing the copyright transfer agreement**

If the OnlineOpen option is not selected the corresponding author will be presented with the copyright transfer agreement (CTA) to sign. The terms and conditions of the CTA can be previewed in the samples associated with the Copyright FAQs below:

CTA Terms and Conditions

[http://authorservices.wiley.com/bauthor/faqs\\_copyright.asp](http://authorservices.wiley.com/bauthor/faqs_copyright.asp)

### **For authors choosing OnlineOpen**

If the OnlineOpen option is selected the corresponding author will have a choice of the following Creative Commons License Open Access Agreements (OAA):

Creative Commons Attribution Non-Commercial License OAA  
Creative Commons Attribution Non-Commercial -NoDerivs License OAA

To preview the terms and conditions of these open access agreements please visit the Copyright FAQs hosted on Wiley Author Services

[http://authorservices.wiley.com/bauthor/faqs\\_copyright.asp](http://authorservices.wiley.com/bauthor/faqs_copyright.asp) and visit

<http://www.wileyopenaccess.com/details/content/12f25db4c87/Copyright--License.html>.

If you select the OnlineOpen option and your research is funded by The Wellcome Trust and members of the Research Councils UK (RCUK) you will be given the opportunity to publish your article under a CC-BY license supporting you in complying with Wellcome Trust and Research Councils UK requirements. For more information on this policy and the Journal's compliant self-archiving policy please visit:

<http://www.wiley.com/go/funderstatement>.

For RCUK and Wellcome Trust authors click on the link below to preview the terms and conditions of this license:

#### Creative Commons Attribution License OAA

To preview the terms and conditions of these open access agreements please visit the Copyright FAQs hosted on Wiley Author Services

[http://authorservices.wiley.com/bauthor/faqs\\_copyright.asp](http://authorservices.wiley.com/bauthor/faqs_copyright.asp) and visit

<http://www.wileyopenaccess.com/details/content/12f25db4c87/Copyright--License.html>.

**Proofs.** Page proofs will be sent to the corresponding author and should be returned within 3 days of receipt to the Production Editor. Significant textual alterations are unacceptable at proof stage without the written approval of the Editor, and they are likely to result in the delay of publication.

The corresponding author will receive an e-mail alert containing a link to a secure web site. A working e-mail address must therefore be provided for the corresponding author. In the absence of the corresponding author, please arrange for a colleague to access the e-mail to retrieve the proofs.

Please note that you have final responsibility for what is stated in the proofs of your manuscript. However, the proofs are checked thoroughly by the *JEADV* editorial team

and if we do not receive corrections after several automated reminders to the email address supplied for the corresponding author then we will assume that we have your approval for publication.

**Author services.** Author Services enables authors to track their article—once it has been accepted—through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The author will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript. Visit <http://authorservices.wiley.com/bauthor> for more details on online production tracking and for a wealth of resources including FAQs and tips on article preparation, submission and more.

**Early View.** *JEADV* is covered by the Publisher's Early View service. Early View articles are complete full-text articles published online in advance of their publication in a printed issue. Articles are therefore available as soon as they are ready, rather than having to wait for the next scheduled print issue. Early View articles are complete and final. They have been fully reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after online publication. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the traditional way. They are therefore given a Digital Object Identifier (DOI), which allows the article to be cited and tracked before it is allocated to an issue. After print publication, the DOI remains valid and can continue to be used to cite and access the article. More information about DOIs can be found online at <http://www.doi.org/faq.html>.

**Offprints.** Free access to the final PDF offprint of your article will be available via author services only. Please therefore sign up for author services if you would like to access your article PDF offprint and enjoy the many other benefits the service offers.

Additional offprints can be ordered on the offprint order form, which accompanies the proofs.

**Disclaimer.** The Publisher, the European Academy of Dermatology and Venereology (EADV) and Editors cannot be held responsible for errors or any consequences arising from the use of information contained in this journal; the views and opinions expressed do not necessarily reflect those of the Publisher, the EADV and Editors or the products advertised.